

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION Unclassified		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution is unlimited	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE		5. MONITORING ORGANIZATION REPORT NUMBER(S)	
4. PERFORMING ORGANIZATION REPORT NUMBER(S)		6a. NAME OF PERFORMING ORGANIZATION Rensselaer Polytechnic Institute	
6c. ADDRESS (City, State, and ZIP Code) Troy, NY 12180-3590		6b. OFFICE SYMBOL (If applicable) Chemical Eng.	7a. NAME OF MONITORING ORGANIZATION ONR, Code 333
8a. NAME OF FUNDING/SPONSORING ORGANIZATION ONR, Code 333		8b. OFFICE SYMBOL (If applicable)	7b. ADDRESS (City, State, and ZIP Code) 800 North Quincy St., Room 504 Arlington, VA 22217
8c. ADDRESS (City, State, and ZIP Code) 800 North Quincy St., Room 504 Arlington, VA 22217		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N-00014-94-1-0097	
11. TITLE (Include Security Classification) Gas-Phase Kinetics Measurements for Underwater Explosives		10. SOURCE OF FUNDING NUMBERS	
12. PERSONAL AUTHOR(S) Arthur Fontijn		PROGRAM ELEMENT NO. 601153N	PROJECT NO. 02402
13a. TYPE OF REPORT Annual	13b. TIME COVERED FROM 10/15/96 TO 10/14/97	14. DATE OF REPORT (Year, Month, Day) 97/10/9	15. PAGE COUNT 4
16. SUPPLEMENTARY NOTATION			

17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	Explosives	Reaction Kinetics
			Fluoro-Nitramino	BO
			Boron	A10

19. ABSTRACT (Continue on reverse if necessary and identify by block number)

To aid in the development of boron-enhanced fluorinitramino explosives, the kinetics of individual reactions are measured over wide temperature ranges. It is found that the pre-exponentials of rate coefficients of BO reactions tend to be two orders of magnitude smaller than for the A1O reactions with the same oxidants. This may explain why no reaction could be observed between BO and CO₂,
 $k < 1 \times 10^{-14} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$.

DRAFT QUALITY INSPECTED 3

20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. Richard S. Miller			22b. TELEPHONE (Include Area Code) (703) 696-4404
			22c. OFFICE SYMBOL 333

PROGRESS

Introduction

The goals of this program are:

- (i) to make accurate measurements, over wide temperature ranges, of rate coefficients for boron combustion in C/H/N/O/F environments, needed for models used in the development of new underwater explosives, and
- (ii) to use the measurements to obtain a further understanding, to allow predictions for additional reactions occurring with such explosives.

To this end measurements are made in unique high temperature reactors of the HTFFR (high-temperature fast-flow reactor) and HTP (high-temperature photochemistry) type. These are to be supplemented, where warranted, by semi-empirical and *ab initio* studies.

Results

A series of measurements have now been made on BO reactions. The remarkable observation is that the pre-exponentials of the rate coefficients are much lower than for the corresponding AlO reactions. Witness the following two pairs, where the rate coefficients are expressed in $\text{cm}^3 \text{molecule}^{-1} \text{s}^{-1}$ units:

- (1) $\text{BO} + \text{O}_2 \rightarrow \text{BO}_2 + \text{O}$; $k(300-1000 \text{ K}) = 7.9 \times 10^{-12} \exp(161 \text{ K/T})$ (Ref. 1)
- (2) $\text{AlO} + \text{O}_2 \rightarrow \text{AlO}_2 + \text{O}$; $k(1200-1690 \text{ K}) = 7.7 \times 10^{-10} \exp(-10008 \text{ K/T})$ (Ref. 2)
- (3) $\text{BO} + \text{HCl} \rightarrow (\text{HBO} + \text{Cl}) \text{ and/or } (\text{OBCl} + \text{H})$; $k(300-760 \text{ K}) = 6.3 \times 10^{-13} \exp(-1403 \text{ K/T})$ (Ref. 1)
- (4) $\text{AlO} + \text{HCl} \rightarrow (\text{HALO} + \text{Cl}) \text{ and/or } (\text{OAlCl} + \text{H})$; $k(440-1590 \text{ K}) = 5.6 \times 10^{-11} \exp(-139 \text{ K/T})$ (Ref. 3)

The exothermic $\text{BO} + \text{CO}_2 \rightarrow \text{BO}_2 + \text{CO}$ reaction (5) was found to be immeasurably slow,¹ $k(300-1200 \text{ K}) < 3 \times 10^{-14}$. The AlO + CO₂ reaction, proceeded with k about 4×10^{-14} in this temperature range.⁴ An *ab initio* study of reaction (5) showed evidence for a BOCO₂ adduct, but no barrier.⁵ Hence, a small pre-exponential should be the cause of the small k(5). Current boron combustion models are very sensitive to this reaction.

PLANS

Further experimental studies planned include the reactions of BO with N₂O, and HF, and of BF with BF₃, O₂ and H₂O.

The reason for the above-noted differences in the pre-exponential parts of the rate coefficients of the monoxides of the two group 13 elements B and Al is being considered. In previous work from this laboratory much smaller differences between BCl and AlCl rate coefficients were observed.^{6,7,8} The BCl reactions with O₂ and CO₂ are actually somewhat faster than their AlCl counterparts. For N₂O the reverse relation holds. Semi-empirical SECI theory was applied successfully to predict the relation between the exponential terms for a series of those reactions.⁹ This approach might be extended to the monoxides. These combined considerations are significant for allowing predictions of k(T) values, in addition to those measured, with greater accuracy than appears in the current literature for such refractory species reactions.

PARTICIPANTS AND CONTACTS

The principal participant on this project was J.-D.R. Rocha, an MS/Ph.D. student.

PUBLICATIONS AND AWARDS

1. D.P. Belyung, G.T. Dalakos, Q. Zhang, J.D.-R. Rocha, and Fontijn A., "Wide Temperature Range Studies of BO and BO₂ Reactions", Abstract for The Fourth International Conference on Chemical Kinetics at NIST, July 1997. A full-length paper by the same title is in preparation.

There were no Awards this year.

REFERENCES

1. D.P. Belyung, G.T. Dalakos, Q. Zhang, J.D.-R. Rocha, and A. Fontijn, "Wide Temperature Range Studies of BO and BO₂ Reactions", Abstract for The Fourth International Conference on Chemical Kinetics, NIST, Gaithersburg, MD, July 1997.
2. D.P. Belyung and A. Fontijn, "The AlO + O₂ Reaction System over a Wide Temperature Range", *J. Phys. Chem.* 99, 12225 (1995).
3. A.G. Slavejkov, C.T. Stanton, and A. Fontijn, "High-Temperature Fast-Flow Reactor Kinetics Studies of the Reactions of AlO with Cl₂ and HCl over Wide-Temperature Ranges", *J. Phys. Chem.* 94, 3347 (1990).
4. D.F. Rogowski, A.J. English, and A. Fontijn, "A High-Temperature Fast-Flow Reactor Kinetics Study of the Reaction AlO + CO₂ → AlO₂ + CO. Thermochemical Implications" *J. Phys. Chem.* 90, 1688 (1986).
5. P. Marshall, University of North Texas, Private Communication.

6. A.G. Slavejkov, P.M. Futerko, and A. Fontijn, "High-Temperature Fast-Flow Reactor Kinetics Study of the Reaction Between BCl and CO₂ from 770 to 1830 K" *Twenty-Third Symposium (International) on Combustion*, (The Combustion Institute, Pittsburgh, 1990), p. 155
7. P.M. Futerko and A. Fontijn, "Experimental and Transition-State Theory Studies of the Gas-Phase Reactions of AlCl with N₂O, CO₂, and SO₂", *J. Phys. Chem.* 97, 7222 (1993).
8. P.M. Futerko, A.G. Slavejkov, and A. Fontijn, "Wide Temperature Range Kinetics of the Gas-Phase Reactions of BCl with SO₂, N₂O, O₂, and CO₂", *J. Phys. Chem.* 97, 11950 (1993).
9. A.S. Blue, D.P. Belyung, and A. Fontijn, "Activation Barriers for Series of Exothermic Homologous Reactions. V. Boron Group Diatomic Species Reactions", *J. Chem. Phys.* 107, 3791 (1997).